

**SOLUTION/ ANSWER KEY OF PRACTICE PAPER -I**

**CLASS XII MATHEMATICS**

**2019-20**

Q NO	VALUE POINTS
1	(D) square matrix
2	(C) $A^2 - B^2 + BA - AB$
3	(D)8
4	(C) 1/2
5	(D) $(\alpha, \beta, -\gamma)$
6	$(B) \frac{2}{5}$
7	(B) 9/10
8	(C) $\tan x - \cot x + c$
9	(A) (2,0,0)
10	(D) $\frac{2}{\sqrt{29}}$ units
11	$R = \{(3,8), (6,6), (9,4), (12,2)\}$
12	$a=2$
13	$X+y=0$  OR  $(-\infty, -1)$
14	$y=2$
15	$\left( \frac{\vec{a} \cdot \vec{b}}{ \vec{b} ^2} \right) \vec{b}$ OR $2/3, 2/3, -1/3$
16	-Interchanging rows and column we get $\Delta = \begin{vmatrix} 0 & a-b & a-c \\ b-a & 0 & b-c \\ c-a & c-b & 0 \end{vmatrix}$  Taking (-1) common from $R_1, R_2, R_3$ we get $\Delta = (-1)^3 \begin{vmatrix} 0 & b-a & c-a \\ a-b & 0 & c-b \\ a-c & b-c & 0 \end{vmatrix} = -\Delta$  therefore $2\Delta=0 \therefore \Delta=0$
17	$x \log x - x + c$
18	4 OR

	$x \tan \frac{x}{2} + c$
19	$e^x \cos x + c$
20	$yx = \frac{x^2}{2} + c$
21	<p>Let <math>\vec{c}</math> denote the sum of <math>\vec{a}</math> &amp; <math>\vec{b}</math> we have <math>\vec{c} = (2\hat{i} - \hat{j} + 2\hat{k}) + (-\hat{i} + \hat{j} + 3\hat{k}) = \hat{i} + 5\hat{k}</math> now  <math> \vec{c}  = \sqrt{1^2 + 2^2} = \sqrt{26}</math>  Required unit vector is <math>\hat{c} = \frac{\vec{c}}{ \vec{c} } = \frac{1}{\sqrt{26}}(\hat{i} + 5\hat{k}) = \frac{1}{\sqrt{26}}\hat{i} + \frac{5}{26}\hat{k}</math>  OR  P(2,3,0) and Q(-1,-2,-4)  <math>\overrightarrow{PQ} = (-1, -2)\hat{i} + (-2, -3)\hat{j} + (-4, -0)\hat{k}</math>  <math>= -3\hat{i} - 5\hat{j} - 4\hat{k}</math>  <math>\therefore</math> Vector joining P and Q given by <math>\overrightarrow{PQ} = -3\hat{i} - 5\hat{j} - 4\hat{k}</math></p>
22	<p><math>a = 1/2</math> not reflexive <math>1/2 \leq 1/2</math> so R is not Reflexive  A=9, b=4, c=2, not transitive  OR  <math>\cos \left[ \sin^{-1} \frac{1}{4} + \sec^{-1} \frac{4}{3} \right] = \cos \left[ \sin^{-1} \frac{1}{4} + \cos^{-1} \frac{4}{3} \right]</math>  <math>= \cos \left( \sin^{-1} \frac{1}{4} \right) \cos \left( \cos^{-1} \frac{3}{4} \right) - \sin \sin^{-1} \frac{1}{4} \sin \cos^{-1} \frac{3}{4}</math>  <math>= \frac{3}{4} \sqrt{1 - \left(\frac{1}{4}\right)^2} - \frac{1}{4} \sqrt{1 - \left(\frac{3}{4}\right)^2}</math>  <math>= \frac{3\sqrt{15} - \sqrt{7}}{16}</math></p>
23	<p><math>y = e^{a \cos^{-1} x} \Rightarrow \frac{dy}{dx} = e^{a \cos^{-1} x} (-a) \frac{-a}{\sqrt{1-x^2}}</math>  Therefore, <math>\sqrt{1-x^2} \frac{dy}{dx} = -ay \dots \dots (1)</math>  Differentiating again w.r.t x, we get <math>\sqrt{1-x^2} \frac{d^2y}{dx^2} - \frac{-x}{\sqrt{1-x^2}} \frac{dy}{dx} = -a \frac{dy}{dx}</math>  <math>\Rightarrow (1-x^2) \frac{d^2y}{dx^2} - x \frac{dy}{dx} = -a \sqrt{1-x^2} \frac{dy}{dx}</math>  <math>= -a(-ay)</math> hence <math>(1-x^2) \frac{d^2y}{dx^2} - x \frac{dy}{dx} - a^2 y = 0</math></p>
24	<p>Given, diameter of the balloon = <math>\frac{3}{2}(2x+1)</math>  <math>\therefore</math> Radius of the balloon = <math>\frac{\text{Diameter}}{2}</math>  <math>= \frac{1}{2} \left[ \frac{3}{2}(2x+1) \right] = \frac{3}{4}(2x+1)</math>  For the volume V, the balloon is given by  <math>V = \frac{4}{3} \pi (\text{radius})^3 = \frac{4}{3} \pi \left[ \frac{3}{4}(2x+1) \right]^3 = \frac{9\pi}{16} (2x+1)^3</math>  For the rate of change of volume, differentiate w.r.t x, we get</p>

	$\frac{dV}{dx} = \frac{9\pi}{16} \times 3(2x+1)^2 \times 2 = \frac{27\pi}{8}(2x+1)^2$ <p>Thus, the rate of change of volume is <math>\frac{27\pi}{8}(2x+1)^2</math>.</p>
25	<p>Given equations of lines are <math>\frac{x}{2} = \frac{y}{2} = \frac{z}{1}</math> and <math>\frac{x-5}{4} = \frac{y-2}{1} = \frac{z-3}{8}</math></p> <p>here direction ratios of two lines are (2,2,1) and (4,1,8)</p> <p>Let <math>\theta</math> be the acute angle between the given lines, then</p> $\cos \theta = \frac{ a_1 a_2 + b_1 b_2 + c_1 c_2 }{\sqrt{a_1^2 + b_1^2 + c_1^2} \sqrt{a_2^2 + b_2^2 + c_2^2}}$ $\cos \theta = \frac{ 2 \times 4 + 2 \times 1 + 1 \times 8 }{\sqrt{2^2 + 2^2 + 1^2} \sqrt{4^2 + 1^2 + 8^2}}$ $= \frac{ 8 + 2 + 8 }{\sqrt{4 + 4 + 1} \sqrt{16 + 1 + 64}}$ $= \frac{18}{\sqrt{9} \sqrt{81}}$ $= \frac{18}{3 \times 9} = \frac{2}{3} \Rightarrow \theta = \cos^{-1} \left( \frac{2}{3} \right)$
26	<p>Given <math>n=6</math> and <math>p = \frac{\text{Number of odd number in one die}}{\text{Total number in one die}} = \frac{3}{6} = \frac{1}{2}</math></p> $\therefore q = 1 - p = 1 - \frac{1}{2}$ <p>So . <math>P(\text{getting a 5 success in six trials}) = P(X=5) = {}^6C_5 p^5 q^1 = 1 \times \left(\frac{1}{2}\right)^5 \left(\frac{1}{2}\right)^1 = 1/64</math></p>
27	<p>(i) Reflexive: <math>\forall a \in A,  a-a =0</math> which is even  <math>\Rightarrow (a,a) \in R</math>, hence <math>R</math> is reflexive.</p> <p>(ii) Symmetric: Let <math>(a,b) \in R</math>  <math>\Rightarrow  a-b </math> is even  <math>\Rightarrow  -(b-a) </math> is even  <math>\Rightarrow  (b-a) </math> is even          So, <math>(b,a) \in R</math>          Hence, <math>R</math> is symmetric.</p> <p>(iii) Transitive: Let <math>(a,b), (b,c) \in R</math>          So, <math> a-b </math> is even and <math> b-c </math> is even  <math>\Rightarrow a-b=2\lambda, b-c=2\mu</math> where <math>\lambda, \mu \in Z</math>          Now, <math>a-c = (a-b) + (b-c) = 2(\lambda + \mu)</math>  <math>\Rightarrow  a-c </math> is even, hence <math>R</math> is transitive.          Since <math>R</math> is reflexive, symmetric, transitive          Therefore, it is an equivalence relation.</p>

28

$$\frac{dy}{dx} = \frac{x+y}{x-y} = \frac{1+\frac{y}{x}}{1-\frac{y}{x}}$$

Put  $y/x = v$  so that  $\frac{dy}{dx} = v + x \frac{dv}{dx}$

$$\therefore v + x \frac{dv}{dx} = \frac{1+v}{1-v} \Rightarrow x \frac{dv}{dx} = \frac{1+v}{1-v} - v = \frac{1+v-v+v^2}{1-v}$$

$$\Rightarrow \int \frac{1-v}{1+v^2} dv = \int \frac{dx}{x} \Rightarrow \int \frac{1}{1+v^2} dv - \frac{1}{2} \int \frac{2v}{1+v^2} dv = \int \frac{dx}{x}$$

$$\Rightarrow \tan^{-1} v = \frac{1}{2} \log |1+v^2| + \log |x| + c$$

$$\Rightarrow \tan^{-1} \left( \frac{y}{x} \right) = \frac{1}{2} \log \left| \frac{x^2 + y^2}{x^2} \right| + \log |x| + c$$

$$\text{or } \tan^{-1} \left( \frac{y}{x} \right) = \frac{1}{2} \log |x^2 + y^2| + c$$

OR

$$(1+x^2)dy + 2xy dx = \cot x dx.$$

$$\Rightarrow \frac{dy}{dx} + \frac{2x}{1+x^2} \cdot y = \frac{\cot x}{1+x^2}$$

$$\text{I.F.} = e^{\int \frac{2x}{1+x^2} dx} = e^{\log(1+x^2)} = (1+x^2)$$

$$\therefore \text{Solution is, } y \cdot (1+x^2) = \int \cot x dx = \log |\sin x| + c$$

$$\text{or } y = \frac{1}{1+x^2} \cdot \log |\sin x| + \frac{c}{1+x^2}$$

29

$$\Rightarrow \frac{dy}{dx} \cdot \log(\cos x) + y(-\tan x) = \log(\sin y) + x \cdot \cot y \cdot \frac{dy}{dx}$$

$$\Rightarrow \frac{dy}{dx} = \frac{y \cdot \tan x + \log(\sin y)}{\log(\cos x) - x \cot y}$$

30

$$\text{RHS} = \int_a^b f(a+b-x) dx = - \int_b^a f(t) dt, \text{ where } a+b-x = t, dx = -dt$$

$$= \int_a^b f(t) dt = \int_a^b f(x) dx = \text{LHS}$$

$$\text{Let } I = \int_{\frac{\pi}{6}}^{\frac{\pi}{3}} \frac{dx}{1+\sqrt{\tan x}} = \int_{\frac{\pi}{6}}^{\frac{\pi}{3}} \frac{\sqrt{\cos x}}{\sqrt{\cos x} + \sqrt{\sin x}} dx \quad \dots(i)$$

$$= \int_{\frac{\pi}{6}}^{\frac{\pi}{3}} \frac{\sqrt{\cos(\pi/2-x)}}{\sqrt{\cos(\pi/2-x)} + \sqrt{\sin(\pi/2-x)}} dx = \int_{\frac{\pi}{6}}^{\frac{\pi}{3}} \frac{\sqrt{\sin x}}{\sqrt{\sin x} + \sqrt{\cos x}} dx \quad \dots(ii)$$

$$\text{adding (i) and (ii) to get } 2I = \int_{\frac{\pi}{6}}^{\frac{\pi}{3}} 1 \cdot dx = x \Big|_{\frac{\pi}{6}}^{\frac{\pi}{3}} = \pi/6.$$

$$\Rightarrow I = \frac{\pi}{12}$$

31

For the first die :  $P(6)=1/2$  ,  $P(6')=1/2$

i.e.,  $P(6')=P(1)+P(2)+P(3)+P(4)+P(5)=1/2$

$\Rightarrow P(1)=1/10$  ,  $P(1')=9/10$  [  $\therefore P(1) = P(2) = P(3) = P(4) = P(5)$  ]

For the second die:  $P(1)=2/5$  ,  $P(1')=3/5$

Let X: number of ones seen  $\therefore X = 0,1,2$

$$P(X=0)=P(\text{not } 1 \text{ from } 1^{\text{st}} \text{ die}) \cdot P(\text{not } 1 \text{ from } 2^{\text{nd}} \text{ die}) = \frac{9}{10} \times \frac{3}{5} = \frac{27}{50} = 0.54$$

$P(X=1) = P(1 \text{ from } 1^{\text{st}} \text{ die}) P(\text{not } 1 \text{ from } 2^{\text{nd}} \text{ die}) + P(\text{not } 1 \text{ from } 1^{\text{st}} \text{ die}) P(1 \text{ from } 2^{\text{nd}} \text{ die})$


$$= \frac{1}{10} \times \frac{3}{5} + \frac{9}{10} \times \frac{2}{5} = \frac{21}{50} = 0.42$$

$$P(X=2) = P(1 \text{ from } 1^{\text{st}} \text{ die}) P(1 \text{ from } 2^{\text{nd}} \text{ die}) = \frac{1}{10} \times \frac{2}{5} = \frac{2}{50} = 0.04$$

The table for probability distribution is shown as below:

X	0	1	2
P(X)	0.54	0.42	0.04

32



Let  $x$  kg of food 1 be mixed with  $y$  kg of food 2 ...  
 To minimize  $Z = ₹ (50x + 70y)$   
 Subject to the constraints :  
 $2x + y \geq 8$ ,  $x + 2y \geq 10$ ,  $x \geq 0$ ,  $y \geq 0$

Corner Points	Value of $Z$ (in ₹)
A(0, 8)	560
B(2, 4)	380 ← Min. value
C(10, 0)	500

Since feasible region is unbounded so, 380 may or may not be minimum value of  $z$ .

To check, draw  $50x + 70y < 380$  i.e.,  $5x + 7y < 38$ .

As in the half plane  $5x + 7y < 38$ , there is no point common with the feasible region.

Hence minimum value of  $Z$  is ₹ 380.

33

$$\text{Area of ellipse} = 4 \left( \frac{b}{a} \int_0^a \sqrt{a^2 - x^2} dx \right)$$

$$= 4 \left[ \left( \frac{b}{a} \left( \frac{x}{2} \sqrt{a^2 - x^2} + \frac{a^2}{2} \sin^{-1} \frac{x}{a} \right) \right) \right]_0^a$$

$$= 4 \frac{b}{a} \left( \frac{\pi a^2}{4} \right)$$

$$= \pi ab$$

OR

$$a = 1, b = 3, nh = 2$$

$$\int_3^3 (x^2 + x + e^x) dx = \lim_{h \rightarrow 0} h(f(1) + f(1+h) + \dots + f(1+(n-1)h))$$

$$= \lim_{h \rightarrow 0} h(2 + e + (1+h)^2 + (1+h) + e^{1+h} + \dots + (1+(n-1)h)^2 + (1+(n-1)h) + e^{1+(n-1)h})$$

$$= \lim_{h \rightarrow 0} h(2 + e + 2 + 3h + h^2 + e^{1+h} + \dots + (1)^2 + (n-1)^2 h^2 + 2(n-1)h + 1 + (n-1)h + e^{1+(n-1)h})$$

34

$$A^3 - 6A^2 + 5A + 11I = O, \text{ Pre-multiplying by } A^{-1}$$

$$\Rightarrow A^2 - 6A + 5I + 11A^{-1} = O \Rightarrow A^{-1} = -\frac{1}{11}(A^2 - 6A + 5I)$$

$$\therefore A^{-1} = \begin{bmatrix} -3/11 & 4/11 & 5/11 \\ 9/11 & -1/11 & -4/11 \\ 5/11 & -3/11 & -1/11 \end{bmatrix}$$

35

As the d.r.'s of parallel lines are proportional so, the equation of line passing through (2, 3, 2) and parallel to  $\vec{r} = (-2\hat{i} + 3\hat{j}) + \lambda(2\hat{i} - 3\hat{j} + 6\hat{k})$  is :  $\vec{r} = (2\hat{i} + 3\hat{j} + 2\hat{k}) + \lambda(2\hat{i} - 3\hat{j} + 6\hat{k})$ .

Now  $\vec{a}_1 = -2\hat{i} + 3\hat{j}$ ,  $\vec{a}_2 = 2\hat{i} + 3\hat{j} + 2\hat{k}$ ,  $\vec{b} = 2\hat{i} - 3\hat{j} + 6\hat{k}$

$$\Rightarrow \vec{a}_2 - \vec{a}_1 = (2\hat{i} + 3\hat{j} + 2\hat{k}) - (-2\hat{i} + 3\hat{j}) = 4\hat{i} + 2\hat{k} \text{ and } (\vec{a}_2 - \vec{a}_1) \times \vec{b} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 4 & 0 & 2 \\ 2 & -3 & 6 \end{vmatrix} = 6\hat{i} - 20\hat{j} - 12\hat{k}$$

$$\therefore \text{S.D.} = \frac{|(\vec{a}_2 - \vec{a}_1) \times \vec{b}|}{|\vec{b}|}$$

$$\Rightarrow = \frac{|6\hat{i} - 20\hat{j} - 12\hat{k}|}{|2\hat{i} - 3\hat{j} + 6\hat{k}|} = \frac{\sqrt{36 + 400 + 144}}{\sqrt{4 + 9 + 36}} = \frac{\sqrt{580}}{7} \text{ Units.}$$

OR

The d.r.'s of normal to the plane are 2, -1, 1.

Since PQ is perpendicular to the plane so, its equation is

$$\frac{x-3}{2} = \frac{y-2}{-1} = \frac{z-1}{1} = \lambda.$$

The coordinates of any random point on the line PQ :

$$Q(2\lambda + 3, -\lambda + 2, \lambda + 1).$$

$\therefore$  Q lies on the plane so,  $2(2\lambda + 3) - (-\lambda + 2) + (\lambda + 1) + 1 = 0$

$$\Rightarrow 6\lambda + 6 = 0 \quad \therefore \lambda = -1$$

$\therefore$  Foot of perpendicular : Q(1, 3, 0).

Distance PQ  $\sqrt{(3-1)^2 + (2-3)^2 + (1-0)^2} = \sqrt{6}$  Units.


Let M( $\alpha, \beta, \gamma$ ) be the image of P in the plane.

So, Q will be mid-point of PM.


$$\text{That is, } Q(1, 3, 0) = Q\left(\frac{\alpha+3}{2}, \frac{\beta+2}{2}, \frac{\gamma+1}{2}\right)$$

On comparing the coordinates, we get:  $\alpha = -1, \beta = 4, \gamma = -1$ .

Therefore, the Image is M (-1, 4, -1).



36



$$\frac{d^2Z}{dx^2} = -12(a+x)x$$

$$\left(\frac{d^2Z}{dx^2}\right)_{x=\frac{a}{2}} = -9a^2 < 0$$

$$CD = \sqrt{a^2 - x^2}$$

$$\text{Area, } A = \frac{1}{2} \times 2\sqrt{a^2 - x^2} (a+x)$$

$$Z = A^2 = (a-x)(a+x)^3$$

$$\frac{dZ}{dx} = 2(a+x)^2(a-2x)$$

$$\frac{dZ}{dx} = 0 \Rightarrow x = \frac{a}{2}$$

$\therefore Z$  is maximum when  $x = \frac{a}{2}$

i.e., Area is maximum when  $x = \frac{a}{2}$

For maximum area

$$\tan \theta = \frac{CD}{AD} = \frac{\sqrt{a^2 - \frac{a^2}{4}}}{a + \frac{a}{2}} = \frac{1}{\sqrt{3}}$$

$$\theta = \frac{\pi}{6}$$